

Zeros of the partition function for a continuum system at first order transitions

Koo-Chul Lee

Department of Physics and the Center for Theoretical Physics

Seoul National University

Seoul, 151-742, Korea

(February 1, 2008)

Abstract

We extend the circle theorem on the zeros of the partition function to a continuum system. We also calculate the exact zeros of the partition function for a finite system where the probability distribution for the order parameter is given by two asymmetric Gaussian peaks. For the temperature driven first order transition in the thermodynamic limit, the locus and the angular density of zeros are given by $r = e^{(\Delta c/2l)\theta^2}$ and $2\pi g(\theta) = l(1 + \frac{3}{2}(\Delta c/l)^2\theta^2)$ respectively in the complex $z(\equiv re^{i\theta})$ -plane where l is the reduced latent heat, Δc is the discontinuity in the reduced specific heat and $z = \exp(1 - T_c/T)$.

02.30.Dk, 02.50.Cw, 05.70.Fh, 64.60.-i

One of the fascinating subjects of equilibrium statistical mechanics is to understand how an analytic partition function acquires a singularity when the system undergoes a phase transition. For the last three decades main focus has been on the second order transition. Only recently a renewed interest in the first order phase transition began to emerge [1].

Since Yang and Lee [2] first published their celebrated papers on the theory of phase transitions and the circle theorem on the zeros of the partition function, there have been many attempts to generalize the theorem [3]. Fisher [4] initiated the study of zeros of the partition function in the complex temperature plane, and Jones [5] proposed a scenario for the first order transition for a continuum system. However very little is known about the distribution of zeros for the continuum case. This is because the partition function for the continuum system is not a polynomial in general and the original proof of the circle theorem relied heavily on particular properties of the coefficients of a polynomial. Recently we have been able to prove the theorem in a quite different approach [6] and this approach allows us to extend the theorem to the continuum case.

We found that the circle theorem follows from a certain mathematical relation which exists between a probability density function and the zeros of its characteristic function. In this paper we first prove three theorems and a corollary without referring to the partition function. When these theorems and the corollary are translated in physical terms we find that, (1) the zeros of the partition function can be expressed in terms of the discontinuities in the derivatives of the free energy across the phase boundary if there is a non-vanishing discontinuity in the first order derivative; (2) there is no zeros in the single phase region where the probability distribution is given by a single Gaussian peak; (3) the zeros of the partition function are calculable exactly at the two phase coexistence point where the probability distribution is given by two asymmetric Gaussian peaks; and (4) the zeros lie on the unit circle if the transition is symmetric.

Furthermore we find the finite size scaling very much similar to that of the discrete system [6]. Therefore this result can again be used, for a continuum system, (1) to resolve the recent controversy over equal weight vs. equal height of the probability distribution

functions [7–9]; and (2) distinguish the first order transition from the second [10,6], just as we have done for the discrete system in ref. [6].

Consider a probability density function $f(x)$ of a random variable X of continuous type, which satisfy i) $f(x) \geq 0$ and ii) $\int_{-\infty}^{\infty} f(x)dx = 1$. The characteristic function [11] of a random variable X is defined by

$$\phi(\omega) = \int_{-\infty}^{\infty} e^{i\omega x} f(x)dx \quad (1)$$

where i is the imaginary unit. The logarithm of $\phi(\omega)$ is known as the second characteristic function or cumulant generating function and denoted by $\psi(\omega)$. That is,

$$\psi(\omega) = \ln(\phi(\omega)) = \sum_{s=1}^{\infty} \gamma_s \frac{(i\omega)^s}{s!}. \quad (2)$$

The expansion coefficients, γ_s are known s -th cumulants or semi-invariants and are calculable by the formula, $\gamma_s = \partial^s \psi(\omega) / \partial (i\omega)^s |_{i\omega=0}$.

If $\phi(\omega)$ can be analytically continued into the complex ω -plane, $\mathcal{M}(t) = \phi(-it)$ is a real function called the moment generating function. Since the characteristic function always exists and its properties are well known [11], we will consider zeros of the characteristic function.

Let S_* be a subset of the sample space of a random variable X . The conditional distribution of X within the sample space, S_* , is called the *truncated distribution* of X , whose probability density function $f_*(x)$ is given by

$$f_*(x) = \frac{f(x)}{\int_{S_*} f(x)dx}, \quad (3)$$

where $x \in S_*$. Suppose the characteristic function of the distribution which can be analytically continued into the complex plane. We will consider zeros of the characteristic function in the finite region only. If the density function is not a delta function (in that case there are no zeros in the finite region) we can always divide the density function into two truncated density functions, taking a point x_* some point in the middle of the distribution.

That is,

$$f(x) = c(f_1(x) + af_2(x)), \quad (4)$$

where

$$f_1(x) = \begin{cases} \frac{f(x)}{\int_{-\infty}^{x_*} f(x)dx} & : x < x_* \\ 0 & : x \geq x_* , \end{cases}$$

$$f_2(x) = \begin{cases} 0 & : x < x_* \\ \frac{f(x)}{\int_{x_*}^{\infty} f(x)dx} & : x \geq x_* , \end{cases}$$

$$c = \int_{-\infty}^{x_*} f(x)dx, \text{ and } a = \int_{x_*}^{\infty} f(x)dx / \int_{-\infty}^{x_*} f(x)dx.$$

Consider the zeros of characteristic function

$$\phi(\omega) = \phi_1(\omega) + a\phi_2(\omega). \quad (5)$$

If $\psi_1(\omega)$ and $\psi_2(\omega)$, the cumulant generating functions, of $\phi_1(\omega)$ and $a\phi_2(\omega)$ exist except at isolated zeros, then we can write eq.(5) in terms of these cumulant generating functions as

$$\phi(\omega) = 2e^{\bar{\psi}} \cosh(\tilde{\psi}(\omega)). \quad (6)$$

In the above $\bar{\psi}(\omega) = (\psi_1(\omega) + \psi_2(\omega))/2$ and $\tilde{\psi}(\omega) = (\psi_2(\omega) - \psi_1(\omega))/2$.

It should be noted that, because of the factor a in $\psi_2(\omega)$, $\psi_2(0) = \ln(a)$. It should further be noted that the zeros of $\phi(\omega)$, in eq.(5) are zeros of $\cosh(\tilde{\psi}(\omega))$ only. This is because any zeros of $e^{\bar{\psi}}$ cancel the poles of $\cosh(\tilde{\psi}(\omega))$. This in turn can be understood because eq.(6) is nothing but $2(a\phi_1\phi_2)^{\frac{1}{2}}[\{\phi_1 + a\phi_2\}/2(a\phi_1\phi_2)^{\frac{1}{2}}]$.

Therefore the zeros of $\phi(\omega)$ may be obtained by solving

$$\tilde{\psi}(\omega) = \pm i(1/2 + k)\pi \equiv iI_k, \quad (7)$$

where $k = 0, 1, 2, \dots$.

Now using the cumulant expansion (2) we can write $\tilde{\psi}(\omega)$ as

$$\tilde{\psi}(\omega) = \sum_{s=1}^{\infty} \tilde{\gamma}_s \frac{(i\omega)^s}{s!}. \quad (8)$$

where $\tilde{\gamma}_s = (\gamma_s^{(2)} - \gamma_s^{(1)})/2$.

If $\tilde{\gamma}_1 = \left[\frac{d}{d(i\omega)} \tilde{\psi}(\omega) \right]_{i\omega=0} \neq 0$, the inverse function $i\omega = \tilde{\psi}^{-1}(\tilde{\psi})$ exists near the origin, and the Taylor series about the origin gives, using the Lagrange's formula [12],

$$i\omega = \sum_{s=1}^{\infty} b_s \frac{\tilde{\psi}^s}{s!}, \quad (9)$$

where

$$b_s = \frac{1}{s!} \frac{d^{s-1}}{d(i\omega)^{s-1}} \left(\frac{i\omega}{\tilde{\psi}(\omega)} \right)^s \Big|_{i\omega=0}. \quad (10)$$

Notice that (9) is the inverted series of (8). It should also be noticed that, since $\tilde{\gamma}_s$'s are real, b_s 's are also real. This leads to theorem I.

Theorem I. The zeros of the characteristic function $\phi(\omega)$ can be expressed by

$$i\omega_k = \sum_{s=1}^{\infty} b_s \frac{(iI_k)^s}{s!}, \quad (11)$$

where b_s 's I_k are given by (10) and (7).

Although the dividing point x_* is arbitrary and b_s 's should not depend on x_* , $\tilde{\gamma}_s$'s have little meaning unless the original probability density function has two separate distributions over two distinct regions and x_* is taken at some point between the two regions.

For a symmetric distribution where $f(x^* + x) = f(x^* - x)$, it is convenient to consider a random variable defined by $Y = X + x^*$. Then the density function becomes symmetric about the origin, $\hat{f}(-y) = \hat{f}(y)$ and the zeros of $\phi_y(\omega)$ are the same as those of $\phi_x(\omega)$ since $\phi_y(\omega) = e^{ix^*\omega} \phi_x(\omega)$.

It can be easily shown that the characteristic function is symmetric about the origin iff the density function is symmetric. Let us take for a symmetric density function $f(y)$ about the origin and the corresponding characteristic functions of truncated distribution functions, $\phi_1(\omega) = 2 \int_{-\infty}^0 e^{i\omega y} f(y) dy$ and $\phi_2(\omega) = 2 \int_0^{\infty} e^{i\omega y} f(y) dy$. Since $\phi_1(-\omega) = \phi_2(\omega)$, we have $\tilde{\gamma}_s = \gamma_s^{(2)} - \gamma_s^{(1)} = 0$, for even s , and $\tilde{\gamma}_s = \gamma_s^{(2)} = -\gamma_s^{(1)}$, for odd s .

Therefore

$$\tilde{\psi}(\omega) = \sum_{s=0}^{\infty} \tilde{\gamma}_{2s+1} \frac{(i\omega)^{2s+1}}{(2s+1)!} = i \sum_{s=0}^{\infty} (-1)^s \tilde{\gamma}_{2s+1} \frac{\omega^{2s+1}}{(2s+1)!}. \quad (12)$$

By the inverse series given by eq.(9), we have

$$\omega = \sum_{s=0}^{\infty} (-1)^s b_{2s+1} \frac{(\tilde{\psi}/i)^{2s+1}}{(2s+1)!}, \quad (13)$$

Substituting the solution for zeros (7) we have

$$\omega_k = \sum_{s=0}^{\infty} (-1)^s b_{2s+1} \frac{I_k^{2s+1}}{(2s+1)!}, \quad (14)$$

provided that series converges. Since b_s 's are real ω_k 's are real. This leads to the following corollary.

Corollary. Zeros of characteristic function of a symmetric distribution function lie on the real axis, provided that the series in (11) converges.

This means that in the complex $z = e^{-i\omega}$ -plane, the Mellin transformation $P(z)$, defined by $P(z) = \phi(-i \ln(z))$ [11] has zeros only on the unit circle.

For some probability distributions we show next that it is possible to calculate zeros of the characteristic function explicitly.

If the density function of the Gaussian distribution is $f(x) = \exp(-\frac{(x-\mu)^2}{2\sigma^2})/\sqrt{2\pi}\sigma$, the cumulant generating function is given by $\psi(\omega) = \mu(i\omega) + \frac{1}{2}\sigma^2(i\omega)^2$ [11]. Since the exponential function of an arbitrary entire function can not have zeros in the finite region [12] and $\psi(\omega)$ in the above is an entire function, $\phi(\omega) = e^{\psi(\omega)}$ can not have zeros in the finite region. This leads to theorem II.

Theorem. II. The characteristic function of the Gaussian distribution has no zeros in the finite region.

On the other hand if the density function of a double Gaussian peak is given by

$$f(x) = e^{-\frac{(x-\mu_1)^2}{2\sigma_1^2}}/\sqrt{2\pi}\sigma_1 + ae^{-\frac{(x-\mu_2)^2}{2\sigma_2^2}}/\sqrt{2\pi}\sigma_2, \quad (15)$$

the cumulant generating functions for the two peaks are given by $\psi_1(\omega) = \mu_1(i\omega) + \frac{1}{2}\sigma_1^2(i\omega)^2$, and $\psi_2(\omega) = \ln(a) + \mu_2(i\omega) + \frac{1}{2}\sigma_2^2(i\omega)^2$ respectively. Therefore $\tilde{\psi}(\omega) = \ln(a)/2 + m(i\omega) + \frac{1}{2}\tilde{\sigma}^2(i\omega)^2$, where $m = (\mu_2 - \mu_1)/2$ and $\tilde{\sigma}^2 = (\sigma_2^2 - \sigma_1^2)/2$. Solving the equation (7) using the above $\tilde{\psi}(\omega)$, the zeros of the characteristic function can be obtained explicitly after performing some algebra. This leads to theorem III.

Theorem. III. If the density function of a double Gaussian peak is given by (15), the zeros of the characteristic function are $\omega_k = \lambda_k/|\tilde{\sigma}| + i(m - |\tilde{\sigma}I_k/\lambda_k|)/\tilde{\sigma}^2$, where $\lambda_k = \pm[\{(\frac{m^2}{2\tilde{\sigma}^2} - \frac{\ln(a)}{2})^2 + I_k^2\}^{\frac{1}{2}} - (\frac{m^2}{2\tilde{\sigma}^2} - \frac{\ln(a)}{2})]^{\frac{1}{2}}$.

In the asymptotic limit where $a \rightarrow 1$ and $\tilde{\sigma} \rightarrow 0$, the zeros are given by

$$\omega_k = I_k/m/(1 - \epsilon_k\tilde{\sigma}^2/m) + i\epsilon_k \quad (16)$$

where $\epsilon_k = \ln(a)/2m - \frac{1}{2}\tilde{\sigma}^2 I_k^2/m^3$. If we put $a = 1$ and $\tilde{\sigma} = 0$ in the above, then we have $\epsilon_k = 0$, which makes ω_k 's real. This is an explicit example of the corollary of theorem I.

The above mathematical results can be readily applied to the theory of the phase transition. As an example consider a canonical partition function $Z(\beta)$ defined by $Z(\beta) = \int_{-\infty}^{\infty} e^{-\beta E} \Omega(E) dE$, where $\Omega(E)$ is the density of states at E , β is the inverse temperature $1/k_b T$ and k_b is the Boltzmann's constant. We can identify this partition function as a moment-generating function $\mathcal{M}(t)$,

$$\mathcal{M}(t) \equiv Z(\beta)/Z(\beta_o) = \int_{-\infty}^{\infty} e^{tx} f(x) dx \quad (17)$$

where the probability density function is given by

$$f(x) = \Omega(x/\beta_o) e^{-x} / \int_{-\infty}^{\infty} \Omega(x/\beta_o) e^{-x} dx. \quad (18)$$

In the above $t = 1 - \beta/\beta_o$, $x = \beta_o E$, and β_o is a reference inverse temperature around which the system fluctuates.

Since the cumulants of $f(x)$ are related to the derivatives of the free energy, theorem I implies that the zeros of the partition function can be expressed in terms of the discontinuities in the derivatives of the free energy provided that the first order derivative has a non-vanishing discontinuity. If $f(x)$ is a symmetric function, then the corollary of theorem I says that the zeros of the partition function lie on the unit circle in the complex e^t -plane. Thus we have extended the Lee-Yang unit circle theorem to the continuum case. From the general principle of statistical mechanics, $f(x)$ can be approximated by the Gaussian distribution. In the thermodynamic limit its mean is given by the internal energy $\beta_o U$, and the variance by the heat capacity as $\sigma^2 = C_v/k_b$. Here β_o is the inverse temperature of a single phase. Due to theorem II, the partition function has no zeros.

On the other hand if the system undergoes a first order transition at $\beta_o = \beta_c$, then $f(x)$ is characterized by a double Gaussian peak [7,8] separated by the discontinuity of the internal energy, or the latent heat in the thermodynamic limit. Let us further assume that the ratio of the weight of the two peaks is a , and there is also a discontinuity in C_v , ΔC_v . Let N be some integer representing an extensive thermodynamic quantity, say the number of particles of the system. Introducing the reduced internal energy $u = \beta_c U/N$, the reduced latent heat $l = \Delta u \equiv u_2 - u_1$ and the reduced specific heat $c = C_v/Nk_b$, theorem III now says that zeros of the partition function for small values of k may be written as,

$$\ln(r_k) = \Re(t_k) = -\ln(a)/Nl + \frac{1}{2}(\Delta c/l)(\vartheta_k/l)^2 \quad (19)$$

$$\theta_k = \Im(t_k) = (\vartheta_k/l) \left\{ 1 + \frac{1}{2}(\Delta c/l) \ln(a)/Nl - \frac{1}{2}(\Delta c/l)^2 (\vartheta_k/l)^2 \right\} \quad (20)$$

Here we have used $\tilde{\sigma}^2 = \Delta c N/2$, $m = lN/2$ and $\vartheta_k = (1 + 2k)\pi/N$ with $k = 0, \pm 1, \pm 2, \dots$. We see that the dominant finite size correction is the terms dependent on the asymmetric factor a .

In the thermodynamic limit where $N \rightarrow \infty$, a dependent terms vanish, and the equation for the locus of zeros becomes

$$r = e^{(\Delta c/2l)\theta^2}, \quad (21)$$

where r_k and θ_k are replaced by the continuous variables r and θ . In Fig. 1, we plot u as a function T/T_c , and zeros in the complex t - and e^t -planes. $l = 1.0$ and $\Delta c = \pm 0.2, 0$ are used in all three figures and the exact zeros given by theorem III are calculated using $N = 20$ and $a = 1$.

The angular density of zeros defined by $Ng(\theta) = 1/(\theta_{k+1} - \theta_k)$ can be written for small values of k , as

$$2\pi g(\theta) = l \left\{ 1 + \frac{3}{2} (\Delta c/l)^2 \theta^2 \right\}. \quad (22)$$

If the transition is symmetric ($\Delta c = 0$), the locus of zeros becomes the unit circle with the uniform density $2\pi g(\theta) = l$. One should note that the equation for the locus of zeros is valid only near the real axis. This is because terms beyond the Gaussian approximation become important as the argument θ grows, *i.e.* for large values of k , as we have shown with an example in ref. [6]. Finally it should be remarked that the number of zeros in this example is infinite in the complex t -plane (Fig. 1(b)). However only a finite number of zeros closes the circle in the complex $z = e^t$ for a finite system if we consider only the first Riemann sheet. In fact for the symmetric case where $\Delta c = 0$, there are exactly N zeros distributed uniformly on the unit circle if we scale the energy of the system by l , so that $l = 1.0$.

In conclusion we have shown that the scenario for the first order phase transition put forward by Yang and Lee [2] is valid in a continuum system. Because we have shown a formal relation between the discontinuities in the derivatives of the free energy and the distribution of zeros of the partition function, it can now be applied to any type of first order phase transition. It can also be extended to a multi-phase coexistence point. In this case, one only needs to consider the multi-dimensional complex space as we have done in ref. [6]. The existence of the formal relation presented in this paper was suspected by the

original proponents of the theorem Lee and Yang themselves. In their 1952 paper [2], they expressed their sentiment in the concluding remark by saying, "...distribution (of zeros) should exhibit such simple regularities One can not escape the feeling that there is a very simple basis underlying the theorem, with much wider application, which still has to be discovered." We believe we have discovered this simple basis.

This work was supported in part by the Ministry of Education, Republic of Korea through a grant to the Research Institute for Basic Sciences, Seoul National University, in part by the Korea Science Foundation through Research Grant to the Center for Theoretical Physics, Seoul National University and in part by S.N.U. Daewoo Research Fund.

REFERENCES

- [1] See e.g., *Dynamics of First Order Phase Transitions*, ed. H.J. Herrmann, W. Janke and F. Karsch, (World Scientific, Singapore, 1992) and references therein.
- [2] C.N. Yang and T.D. Lee, Phys. Rev. **87**, 404 (1952); T.D. Lee and C.N. Yang, Phys. Rev. **87**, 410 (1952)
- [3] See for example, R. B. Griffiths in *Phase Transitions and Critical Phenomena* vol. 1 ed. C. Domb and M.S. Green (Academic, New York, 1973) p 8 and references therein.
- [4] M. E. Fisher, in *Lectures in Theoretical Physics* (University of Colorado Press, Boulder, 1965), Vol. 7c
- [5] G.L. Jones J. Math. Phys. **7**, 2000 (1966)
- [6] K-C. Lee, Phys. Rev. Lett. (to be published).
- [7] K. Binder and D.P. Landau, Phys. Rev. **B 30**, 1477 (1984)
- [8] M.S. Challa, D.P. Landau, and K. Binder, Phys. Rev. **B 34**, 1481 (1986)
- [9] C. Borgs and R. Kotecky, J. of Stat. Phys. **61**, 79 (1990)
- [10] K-C. Lee, Phys. Rev. E **48** 3459 (1993)
- [11] B.R. Bhat *Modern Probability Theory, 2nd Ed.* (Halsted Press(John Wiley & Sons) 1981); M. Ochi *Applied Probability & Stochastic Processes* (John Wiley & Sons, 1992)
- [12] G.F. Carrier, M. Krook and C.E. Pearson *Functions of Complex Variable* (McGraw-Hill Book Company, 1966) p.62

FIGURES

FIG. 1. (a) u as a function of $T/T_c = 1/(1 - t)$. $u_1 = 0.7$ and $c_1 = 1.0$ are taken arbitrarily. (b) Zeros in the complex t -plane given by theorem III. For $\Delta c > 0$, the line of zeros arches toward the positive real axis. Only zeros between the two dashed lines appear in the first Riemann sheet in the complex e^t -plane. (c) Zeros in the complex e^t -plane. The solid lines are the loci of the zeros in the thermodynamic limit given by (21). For $\Delta c > 0$, the locus lies outside the unit circle which corresponds to the symmetric case, $\Delta c = 0$.